

REVIEW

Shared Pathways: Death Receptors and Cytotoxic Drugs in Cancer Therapy*

Istvan PETAK, Janet A HOUGHTON

Division of Experimental Hematology, Department of Hematology-Oncology, St. Jude Children's Research Hospital, Memphis, TN, USA

Death ligands (TNF, FasL, TRAIL) and their respective death receptor signaling pathways can be used to induce tumor cells to undergo apoptosis. Chemotherapeutic drugs can induce apoptosis and the upregulation of death ligands or their receptors. Downstream events following cytotoxic stress-induced DNA damage and the signaling pathways that lead to the induction of apoptosis may be either dependent or independent of death receptor signaling. The involvement of the Fas signaling pathway in chemotherapy-induced apoptosis has been the most extensively studied, with the current emergence of information on the TRAIL signaling

Keywords: apoptosis, tumor, death receptor, Fas, drug, p53

pathway. Fas-mediated and chemotherapy-induced apoptosis can converge at the level of the receptor, FasL, DISC formation, activation of the initiator caspase-8, at the level of the mitochondria, or at the level of downstream effector caspase activation. Convergence is influenced by the specific form of DNA damage, the cellular environment, and the specific pathway(s) by which death receptor-mediated or drug-mediated apoptosis are induced. This review discusses the different levels of interaction between signaling pathways in the different forms of cell death. (Pathology Oncology Research Vol 7, No 2, 95–106, 2001)

Death receptors and their ligands

Cells of higher metazoans activate their self-destructive mechanisms in response not only to environmental stress, internal damage or lack of survival signals but also by the "instruction" of specific death signals (reviewed in 1,2). Death receptors are cell surface receptors that transmit apoptosis signals initiated by their specific "death ligands". Death receptors belong to the TNF receptor superfamily,

and are defined by homologous cysteine-rich extracellular domains. There are six members of this family (DR1-DR6) that contain a cytoplasmic "death domain". These death domains enable the receptors to deliver apoptotic signals in sensitive cells upon binding of the specific ligand. In addition, there are three "decoy" receptors with functional extracellular domains and absent or truncated death domains that bind death ligands (DcR 1-3), but are unable to transmit a cell death signal (*Figure 1*).

Death ligands

Death ligands that bind to specific receptors of the TNF receptor superfamily are type II transmembrane glycoproteins with an extracellular C-terminus, a retained transmembrane region, and a cytoplasmic tail (reviewed in 1,2)

Death ligands form oligomers, most often trimers, that are important for their ability to crosslink their receptors. Proteolytic processing is a common postranslational feature of death ligands. TNF³ and FasL^{4,5} are each processed by metalloproteinases, whereas TRAIL⁶ is cleaved by cysteine proteases at the cell surface to yield a soluble

Received: May 22, 2001; *accepted:* June 5, 2001

Correspondence: Janet A HOUGHTON; Division of Experimental Hematology, Department of Hematology-Oncology, St. Jude Children's Research Hospital, 332 North Lauderdale, Memphis, TN 38105, USA; Tel: (901)495-3456, fax: (901)523-2622, e-mail: janet.houghton@stjude.org

* This research was supported by NIH Awards RO1 CA 32613, the Cancer Center Support (CORE) Grant CA 21765, and by the American Lebanese Syrian Associated Charities. I. Petak was supported in part by a scholarship from the Fulbright Program, and by the 1st Institute of Pathology and Experimental Cancer Research, Semmelweis University, Budapest, Hungary




Figure 1. Death receptors belong to the TNF receptor superfamily and share a distinctive intracytoplasmic domain called the "death domain". Their respective death ligands can also bind to "decoy" receptors, where the death domain is truncated or absent.

cytokine. Data indicate tissue-specific differences in death ligand processing determined by different expression of various proteases (e.g. MMPs) and protease inhibitors (e.g. TIMPs). Therefore, the posttranslational modification of death ligands may influence their biological activity in different tissues. The membrane-bound position may confer distinct receptor binding and cellular responses not seen in soluble counterparts. Soluble FasL may induce autocrine signaling in the same cell or paracrine signaling to induce apoptosis in neighboring cells.⁷ Others have found that soluble FasL is less potent than the membrane-bound form, and that cell surface "shedding" of the ligand is the mechanism of inactivation of its cytotoxicity.⁴ In xenograft models FasL-expressing tumors attracted infiltrating neutrophil granulocytes resulting in decreased tumor growth compared to FasL-negative tumors. Further, adenoviral mediated gene transfer of FasL induced regression of Fas⁻ CT26 colon carcinoma xenografts, which was mediated by inflammatory cells.⁸

The study of these death receptor/death ligand systems and their signaling mechanisms can influence the search for improved cancer therapy in several ways: 1) Death ligands can be used directly to target cancer cells expressing their specific receptor, 2) DNA damaging agents may induce apoptosis dependent on death receptor signaling, 3) Chemotherapeutic agents and cytokines can modulate the death receptor sensitivity of tumor cells, and 4) Drugs can directly activate or inhibit signal components downstream of death receptors.

Receptors for FasL (Fas [DR2], DcR3)

The cell surface receptor Fas (APO-1, CD95) and its ligand (FasL) regulate apoptosis in cells of the immune system and in neoplastic cells derived from other tissues. However analyses of tissues from mice have demonstrated expression of Fas in tissues largely characterized by

high rates of cell turnover and apoptotic cell death, including epithelial tissues.^{9,10} The decoy receptor DcR3 can also bind FasL and is frequently expressed in malignancies¹¹ but does not necessarily correlate with resistance to Fas-mediated apoptosis. Following trimerization of Fas after ligation, apoptosis is initiated.¹² Several proteins have been identified that bind to the intracellular death domain of Fas to form the death-inducing signaling complex (DISC). The adaptor protein FADD^{13,14} binds to Fas via the death domain, and subsequently recruits procaspase-8 to the DISC.¹⁵⁻¹⁷ Following release of active caspase-8 from the DISC, a cascade of ICE-like proteases is activated. An anti-apoptotic factor, c-FLIP, has been shown to exist in two isoforms, c-FLIP_S and c-FLIP_L, which appears to be the predominant form expressed in mammalian cells.^{18,19} C-FLIP can be recruited to the DISC and can block further recruitment of procaspase-8 into the complex, thereby inhibiting the activation of caspase-8.^{18,20}

In Fas-mediated apoptosis, two signaling pathways have been demonstrated (Figure 2).²⁰⁻²¹ In type I cells, ligation of Fas leads to strong caspase-8 activation at the DISC thereby directly activating other caspases including caspase-3 in the absence of mitochondrial involvement. In type II Fas-mediated cell death, only a small amount of DISC is formed leading to the activation of a small amount caspase-8, which cleaves the cytosolic substrate Bid. The proteolytically modified Bid induces conformational changes in Bax that lead to pore formation in the mitochondrial membrane. This process can be blocked by anti-apoptotic members of the Bcl-2 family.²² Release of mitochondrial cytochrome c and dATP triggers the formation of the "apoptosome" with caspase-9 and the adaptor molecule Apaf-1. In the apoptosome caspase-9 undergoes autoactivation and further activates the effector caspases-3, -6, and -7.²³ The active caspase-6 can activate additional caspase-8 molecules such that the apoptotic signal of Fas can be further amplified.^{21,22}

Receptors for TRAIL (DR4, DR5, DcR1, DcR2)

In contrast to the Fas signaling pathway, considerably less is known about the biologic function of TRAIL in the regulation of apoptosis in cells and tissues. Also, in contrast to the Fas system, the only primary cells identified as susceptible to apoptosis induction by TRAIL are activated T lymphocytes, suggesting a role in limiting the immune response,^{24,25} hence normal cells are in general TRAIL-resistant. In contrast, tumor cell lines of diverse origin are sensitive to TRAIL.^{6,26-29} Like FasL, TRAIL is a type II transmembrane protein with an intracellular amino terminal portion and extracellular carboxyl terminus, and functions optimally in the trimeric form.^{28,29} In contrast to other members of the TNF family whose expression is tightly

regulated and only transiently expressed on activated cells, mRNA for TRAIL is detected in a wide range of tissues including peripheral blood lymphocytes, spleen, thymus,




Figure 2. Induction of apoptosis by death ligands and chemotherapeutic drugs: Upon activation of death receptors by their ligands a death-inducing signaling complex (DISC) is formed at the receptor. Activated caspase-8 is released and can directly activate caspase-3 in type I cells. Alternatively, further amplification of the apoptotic signal is required with the involvement of the mitochondria in type II cells. Cleavage of Bid by caspase-8 triggers the release of cytochrome C and Smac/DIABLO from the mitochondria. Cytochrome C activates formation of the apoptosome and subsequently the activation of caspase-9. Smac/DIABLO facilitates the activation of caspase-3 and caspase-9 by blocking the caspase inhibitory proteins (IAP1/2). The activation of caspase-6 can lead to the further activation of caspase-8. Cytotoxic drugs can directly target the mitochondria or can activate various signaling pathways. P53 can induce apoptosis by translocating to the mitochondria, by increasing the expression of the death receptors or pro-apoptotic Bax protein, or by production of reactive oxidative species (ROS). ROS activates the transcription factor NF-kB, which increases the expression of several death ligands including FasL. Stress response signals to drugs include the jun kinase/stress activated protein kinase (JNK/SAPK) system, leading to the activation of AP-1 which transcriptionally regulates the expression of death ligands. Increased production of ceramide can indirectly induce the mitochondrial release of pro-apoptotic molecules and also the inhibition of the anti-apoptotic Akt/PKB kinase. Drugs can also inhibit the expression of the anti-apoptotic c-FLIP molecule and may induce direct activation of receptor aggregation, DISC formation, and caspase-8 activation by unknown mechanisms.

prostate, ovary, colon and placenta, but not brain, liver or testis.³⁰ There are four receptors for TRAIL. DR4³¹ and DR5,^{32,33} like Fas, are type I transmembrane proteins with extracellular cysteine-rich residues and a cytoplasmic death domain. There are two non-death-domain-containing TRAIL receptors, DcR1 (lacking a cytoplasmic domain;³⁴) and DcR2 (containing a truncated death domain;^{35,36}), that bind the ligand with comparable affinity to DR4 and DR5 but cannot transmit a cell death signal, and may protect normal tissues from the cytotoxic action of TRAIL, providing a rationale for the use of TRAIL as a cytotoxic. In neoplastic diseases, DR4 and DR5 are expressed in primary human brain tumors,³⁷ melanomas³⁸ including those resistant to FasL,³⁹ and colon carcinomas.^{28,40} Further, TRAIL has induced apoptosis in malignant glioma cell lines,⁴¹ melanoma cell lines,³⁸ breast and colon carcinoma cell lines.^{28,40}

C-FLIP can also inhibit TRAIL-induced apoptosis.^{38,42} To date, little information is available concerning the receptor complexes for the TRAIL receptors DR4 and DR5. Studies have reported direct binding of the adaptor proteins FADD and TRADD to these receptors and inhibition of TRAIL-induced apoptosis in the presence of dominant negative (DN) mutant forms of FADD and TRADD.³⁸ Further, TRAIL-induced DISC formation has been demonstrated with recruitment of FADD and procaspase-8, similar to the Fas-DISC.⁴²⁻⁴⁵ Hence TRAIL signals through formation of a DISC downstream of DR4 or DR5, which explains an inhibitory role for c-FLIP in TRAIL-induced apoptosis.

Apoptosis induced by chemotherapy

It is known that tumor cells can die in response to sub-lethal stimuli, such that reversible damage can trigger apoptosis. However, when irreversible lethal damage occurs, cell death is not necessarily dependent on the caspase-mediated apoptotic machinery. Irreversible damage can cause cell death without initiating apoptosis, or if triggered, cell death is still not necessarily dependent on caspase activation. Thus, discrepancies can exist between apoptosis, growth inhibition and clonogenic assays.

One of the major goals in cancer therapy is to trigger tumor selective cell death. The discovery of distinct, highly efficient molecular pathways that link cellular stress to cell death led to an explosion of apoptosis research, and its relevance to cancer chemotherapy. It is now widely accepted that in tumor cells many anticancer agents act as apoptosis inducers, and an important determinant of drug resistance is the inability of drugs to trigger apoptosis.^{46,47} Although the primary intracellular targets of action of chemotherapeutic agents are distinct, it has become evident that induced cytotoxicity ultimately converges on a common pathway that induces apopto-

sis.⁴⁸ Cells treated with cytotoxic agents can show the typical characteristics of apoptosis analogous to apoptosis induced by “physiologic” stimuli such as growth factor withdrawal or death receptor activation. Anti-apoptosis genes including Bcl-2 and Bcl-xL can inhibit apoptosis induced by a wide variety of stimuli including chemotherapeutic drugs.⁴⁹ Further, the p53 tumor suppressor gene plays an important role in the regulation of apoptosis induced by DNA damage, which can influence tumor sensitivity to cancer therapy.⁵⁰ Most importantly, drug-induced cell death is frequently mediated by caspase-dependent apoptosis.⁵¹

For hematologic malignancies and certain drug-sensitive solid tumors there is rapid induction of apoptosis following drug treatment, where apoptosis has proven to be a prominent mechanism associated with the induction of tumor remission. In contrast, in other solid malignancies, the association is less clear (Reviewed in 48). Hematopoietic cells can undergo rapid apoptosis within a few hours,⁵² while epithelial cells require at least 24 hours,⁵³ hence the form and kinetics of cell death are clearly dependent upon the cell type. Further, the drug concentration not only influences the cytotoxic response in a quantitative manner, but can also trigger qualitatively different response mechanisms.^{54,57} For doxorubicin and daunomycin, EC₉₀ concentrations led to G2/M arrest and cytostasis in HeLa cells in the absence of inhibition of DNA synthesis, ultimately resulting in delayed apoptosis and cell death. In contrast at 10 x EC₉₀ concentrations, apoptosis was detected as early as 3 hours after drug treatment.⁵⁵ The delay or inhibition of an apoptotic response does not always correlate with long-term cell survival. In HeLa cells rapid apoptosis induced by VP-16 was prevented by Bcl-2, while the loss in clonogenic survival and the formation of giant, multinucleated cells characteristic of mitotic catastrophe, was not affected by Bcl-2.⁵⁷ It has been demonstrated that caspase inhibitors can inhibit apoptosis induced by VP-16 in colon carcinoma cells without affecting loss in clonogenic survival.⁵³ In contrast, DNA damage induced by *5-fluorouracil/leucovorin* (FUra/LV) triggered apoptosis and loss in clonogenic potential, and both were prevented by inhibition of caspase activation.⁵³ Chemotherapeutic agents can induce loss in clonogenic survival at therapeutically relevant concentrations without acute induction, but with delayed induction, of apoptosis.⁵³

Although many different anticancer drugs ultimately mediate cell death by activating the caspase signaling cascade, the molecular mechanisms linking the primary drug target and diverse cellular responses to apoptotic effectors are unclear. The present discussion will examine how drug-induced DNA damage activates apoptosis, and how this may be involved or converge on death receptor-mediated apoptosis pathways.

Sharing pathways between drug – and death receptor – induced apoptosis

Some reports indicate that certain DNA damaging agents may induce apoptosis via Fas signaling,^{58,61} while others suggest that stress-induced apoptosis caused by drug treatment may be independent of Fas^{62,63} (*Figure 2*). Fas dependency of apoptosis can require interaction between Fas and FasL. However other possibilities exist for crosstalk between pathways of drug-induced cell death and death receptor-induced apoptosis. It is clear that drugs and death receptor ligands induce apoptosis that can converge at the level of the death receptor itself, the initiation phase of caspase activation, or via activation of the same effector caspases resulting in similar morphologic and biochemical features of cell death. Evidence is also emerging that chemotherapeutic agent-induced signaling and signaling via the TRAIL receptors can be convergent in the induction of apoptosis.^{64,66}

Sharing the death receptors

In Jurkat T leukemia cells, apoptosis induced by doxorubicin appeared to proceed via activation of FasL,⁵⁸ analogous to the dependence of activation-induced apoptosis in T lymphocytes upon transcription and expression of FasL.^{67,68} In addition, competitive binding of either FasL by using a Fas-Fc chimeric fusion protein,⁵⁸ or of Fas by using a neutralizing anti-Fas Ab, effectively inhibited death induced by VP-16 and VM-26,^{60,61} and the early stages of doxorubicin-induced apoptosis,⁶⁹ suggesting that apoptosis induced by these agents was mediated via Fas signaling. In Type I cells (BJAB), both the receptor and the mitochondrial pathway were activated upon treatment with doxorubicin or VP-16, since blockade of the receptor pathway by overexpression of DN-FADD or of the mitochondrial pathway by Bcl-xL only partially inhibited apoptosis.⁷⁰ Also in these cells, drug treatment induced formation of a FADD- and caspase-8-containing Fas-DISC. It is also well documented that thymineless stress-induced apoptosis induced in thymidylate synthase-deficient (TS⁻) human colon carcinoma cells,⁷¹ wt colon carcinoma cells⁷² and other cell types^{73,74} treated with FUra/LV, that apoptosis signals via Fas both in vitro⁷¹⁻⁷³ and in vivo.⁷⁴ Sensitization of colon carcinoma cells to FUra/LV has been enhanced following upregulated expression of Fas by the cytokine IFN- γ .⁷² 5-Fluorouracil and other cytotoxic agents upregulate the expression of FasL,^{58,71-73,75} which can be transcriptionally regulated by NF- κ B^{58,75} or AP-1,^{58,73,75} and also upregulate Fas, dependent upon p53.^{76,77} In some systems however from data derived using blocking Abs against Fas,^{63,78,80} FasL,⁵³ or DN-FADD,^{80,81} drug-induced apoptosis clearly proceeds via a Fas-independent pathway. This has also been demonstrated in cells

selected for resistance to Fas-mediated apoptosis.^{53,63,82} Interaction of anticancer drugs with the TRAIL receptors is due to upregulated expression of DR5 which enhances the induction of apoptosis.^{61,65}

Sharing the caspase signaling cascade

Following initial caspase activation, these molecules subsequently activate each other in the signaling cascade leading to apoptosis, and form positive feedback loops. There may be a critical threshold level of caspase activation for the "chain-reaction" to be initiated (e.g. Type I vs. Type II cells in Fas-mediated apoptosis). It is possible that independent mechanisms can synergistically activate the same caspases, whether at the level of initiator caspases (e.g. caspase-8) or effector caspases (e.g. caspase-3). This mechanism is a general way of positive interaction between death receptor-induced and drug-induced cell death. Drug-induced and death receptor-induced apoptosis pathways can converge upstream of the mitochondria at the level of caspase-8, or downstream of the mitochondria at the level of effector caspases. Further, drug-induced downregulation of inhibitory factors (c-FLIP, Bcl-2) also enhance caspase activation.

A. Upstream of the mitochondria

It has been demonstrated that anticancer drugs can activate caspase-8 dependent upon or independent of signaling via Fas. Doxorubicin or VP-16 activated caspase-8 as the most apical caspase in Type I cells (BJAB) downstream of DISC formation.⁷⁰ Bid was cleaved prior to changes at the level of the mitochondria demonstrating activation of caspase-8 upstream of these organelles. This has also been reported for staurosporine-mediated apoptosis.⁸³ In addition doxorubicin-induced apoptosis involved FasL-mediated aggregation of Fas followed by DISC formation and activation of caspase-8 in H9 T leukemia cells.⁶⁹ However other reports have demonstrated initial caspase-8 activation independent of Fas.^{82,84} In the NSCLC cell line H460, overexpression of DN-FADD did not prevent procaspase-8 cleavage and subsequent apoptosis induced by cisplatin, topotecan or gemcitabine.⁸⁴ Further overexpression of caspase-9S or XIAP, both inhibitors of caspase-9 activation, failed to block drug-induced apoptosis, indicating lack of mitochondrial involvement in the apoptotic response. In the Jurkat T leukemia model, cells resistant to Fas-mediated apoptosis were equally susceptible to anticancer drugs (doxorubicin, daunorubicin, VP-16, mitomycin-C) and activated caspase-8 with similar kinetics and dose response as Fas-sensitive cells.⁸² In addition a neutralizing Fas decoy or DN-FADD selectively abrogated Fas but not drug-induced effects.

Another level upstream of the mitochondria at which Fas-induced and drug-induced apoptosis may converge is at the

level of c-FLIP expression, which may determine caspase-8 activation, dependent upon death receptor signaling. In human osteosarcoma MG-63 cells cisplatin downregulated expression of c-FLIP_I in a time-dependent manner thereby sensitizing the cells to Fas-mediated apoptosis, but did not influence the expression of other anti-apoptotic proteins including XIAP, c-IAP-1 or c-IAP-2.⁸⁵ Down regulated expression of c-FLIP_L using antisense oligonucleotides sensitized MG-63 cells to Fas-mediated apoptosis thereby confirming the importance of c-FLIP in the mechanism of sensitization. Inhibitors of protein (cycloheximide) or RNA (actinomycin-D) synthesis have also downregulated c-FLIP expression and sensitized cancer cells to Fas-mediated apoptosis^{86,87} and TRAIL-induced apoptosis.^{88,89} However this is not a universal mechanism in all cell types, since Jurkat cells transfected with c-FLIP remained sensitive to doxorubicin, VP-16 and vincristine.⁹⁰

B. Downstream of the mitochondria

Depolarization of the mitochondrial membrane and downstream activation of effector caspases is part of the apoptotic machinery in both drug- and death receptor-induced apoptosis. Deficient activation of caspases leads to cross resistance between these two types of induced cell death,⁹¹ and hence the effector caspases form an additional level at which these two signaling pathways can converge. In Type II Fas-mediated apoptosis caspase-8 cleavage occurs predominantly downstream of the mitochondria in contrast to Type I.²⁰ In Jurkat cells (Type II), variant clones selected for resistance to Fas-mediated apoptosis were cross resistant to VP-16.⁹² The pathways did not converge at the level of Fas ligation or caspase-8 signaling, since both effector caspase processing and cytochrome c release were inhibited in the Fas-resistant variants as well as in Bcl-2 transfectants, suggesting the two apoptosis signaling pathways are under common mitochondrial control. The time dependence of activation of caspases-3, -7, -8 and -9 were also similar.⁹³ In B lymphoma cells epirubicin-induced caspase-8 activation occurred independent of death receptor (Fas) signaling and was mediated downstream of caspase-3 activation.⁸¹ Caspase-3 was also activated after treatment with doxorubicin alone or combined with anti-Fas.⁹⁴ These data indicate that drug-induced apoptosis can also act solely in the execution phase of apoptosis, thereby converging downstream of the mitochondria with death receptor signaling pathways.

Convergence at the level of p53

Chemotherapeutic agents including cisplatin, mitomycin, methotrexate, mitoxantrone, doxorubicin and bleomycin can induce Fas expression in a variety of different cell lines, dependent on the presence of a wtp53

gene.^{76,77} The upregulation of Fas by chemotherapeutic agents can also sensitize tumor cells to Fas-mediated apoptosis. This has been substantiated in studies in which the wtp53 gene has been transfected, followed by the induction of Fas and enhancement of apoptosis.⁹⁵⁻⁹⁸ Overexpression of Fas has sensitized cells to p53-mediated apoptosis.⁹⁷ Further, FUra combined with LV upregulated expression of Fas in human colon carcinoma cells, dependent on the presence of a wtp53 gene, and HCT116 cells became sensitive to anti-Fas after FUra/LV treatment.⁷⁷ This in turn may make cancer cells more vulnerable to cytotoxic lymphocytes expressing FasL. Theoretically, a combination of drugs and recombinant FasL or agonistic anti-Fas mAb could have synergistic anti-tumor activity in vivo. However FasL is toxic, primarily due to the induction of apoptosis in the liver.⁹⁹ The death receptor for TRAIL, DR5, is also upregulated by cytotoxic drugs in a p53-dependent manner.¹⁰⁰ In glioma cell lines, cisplatin and VP-16 upregulated DR5 expression, and demonstrated synergistic activity with TRAIL.¹⁰¹ TRAIL has not demonstrated toxicity in vivo and clinical evaluation is currently being initiated using TRAIL as a cytotoxic ligand alone and in combination with chemotherapeutic agents.

Sharing JNK/SAPK and NF- κ B signaling pathways

The most controversial interaction between drugs and death receptor signaling is at the level of upregulation of death ligands. It has been shown in several cell types that FasL,^{58,59} TRAIL¹⁰² and TNF¹⁰² are upregulated in cytotoxic stress induced by anticancer drugs, independent of p53. Upregulation of FasL in colon carcinoma cell lines has been demonstrated in response to thymineless stress, and after treatment with FUra, doxorubicin, VP-16 or topotecan.^{53,71} The transcriptional regulation of FasL is not fully understood. However, there are reports indicating that drugs may activate JNK/SAPK (JNK: c-Jun N-terminal kinases, SAPK: stress-activated protein kinases) and NF- κ B pathways by generating ROS¹⁰³ or via ceramide¹⁰¹ signaling.

Transcription factors including NF- κ B and AP-1 have been implicated in the induction of apoptosis in response to the cytotoxic agents VP-16 and VM-26,⁵⁸ daunorubicin and mitoxantrone,¹⁰⁵ and to ionizing radiation.⁵⁸ NF- κ B is a dimer of two subunits, NFKB1 (p50) and RelA (p65).^{106,107} Activation of NF- κ B is controlled by phosphorylation and proteolysis of an inhibitory subunit, I κ B, which retains the NFKB1/RelA dimer in the cytoplasm. Upon stimulation, I κ B is phosphorylated by an as yet unknown kinase and subsequently degraded.¹⁰⁸ The free NF- κ B dimer then migrates to the nucleus and binds to specific sequences. Such κ B elements are found in promoter regions of genes that are crucial for immune or acute phase responses, including FasL.⁵⁸ AP-1 is a

sequence-specific transcriptional activator composed of members of the Jun and Fos families (reviewed in 109). These proteins associate to form a variety of homo- and hetero- dimers that bind to a common site. A putative AP-1 binding domain has also been identified in the promoter region of FasL.^{58,73} Further implication for the involvement of AP-1 in the induction of apoptosis comes from studies of the involvement of the JNK/SAPK in apoptosis induction. The JNK, members of the MAPK family, phosphorylate the stimulatory sites of c-Jun,¹⁰⁹ and it is well established that AP-1 is induced via activation of the JNK/SAPK pathway.^{110,111} Recent studies have identified JNK to be involved in cellular responses to environmental stresses including uv light,^{112,113} ionizing radiation,^{113,114} DNA-damaging drugs,^{59,115-118} and Fas-mediated¹¹⁹ apoptosis. However, in individual systems, the role of JNK in mediating apoptosis appears dependent upon the agent used to induce DNA damage.^{59,120} Thus, activation of JNK by cytotoxic drugs^{59,118} has led to apoptosis, and dominant negative mutants of some of the elements of this pathway have delayed or significantly inhibited apoptosis induced by certain cytotoxic agents.^{59,121} Ectopic expression of JNK has also induced apoptosis.¹¹³ Additionally, introduction of a mutation into the putative binding site for AP-1 in the promoter region of a FasL reporter construct, resulted in the inhibition of transcriptional activation following VP-16 or VM26 treatment in Jurkat cells.⁵⁸ Similar results were obtained with a FasL reporter construct following mutation of the NF- κ B binding site.⁵⁸ The pathways responsible for NF- κ B activation are considerably less clear, and may or may not involve the JNK/SAPK pathway.^{58,59,122} However, anticancer drugs including VP-16,^{58,123,124} VM-26,⁵⁸ actinomycin-D,¹²⁴ anthracyclines,¹⁰⁵ topoisomerase I inhibitors,¹²⁴ and ara-C^{123,125} have been reported to activate NF- κ B, and inhibition of this activation can inhibit apoptosis.^{58,123}

Sharing the ceramide pathway

Ceramide produced by sphingomyelinases (Smases) has been recognized as an important evolutionary conserved intracellular second messenger.¹²⁶ Sphingomyelin breakdown products have been implicated in both cytotoxic stress (chemotherapy, ionizing radiation, UV, heat, oxidative stress)¹²⁷⁻¹³¹ and death receptor induced apoptosis.¹³² The Smases are sphingomyelin-specific C phospholipases and are identified by their activation requirements such as pH optimum and cation dependence.¹²⁶ A synthetic production of ceramide de novo also occurs via the enzyme ceramide synthase, which in mammalian cells catalyzes the reaction between the sphingoid base sphinganine and fatty acyl-CoA to produce dihydroceramide.¹³³ Ceramide synthase activity has been detected in the endoplasmic reticulum and in the mitochondria.^{134, 135}

Ceramide generation in response to TNFR and Fas activation has been associated with the adaptor molecules TRADD and FADD. Overexpression of these molecules enhanced TNF-induced stimulation of the acidic form of Smase. Ceramide production was shown to be initiator caspase dependent but upstream of effector caspase activation. However, this initiator caspase has yet to be identified distinct of caspase-8 or caspase-10.¹³⁶⁻¹³⁸ In type II Jurkat cells activation of Fas induced elevation in sphingosine and ceramide concentrations.¹³⁹ Both ceramide- and sphingosine- induced apoptosis was dependent on mitochondrial membrane depolarization and the release of cytochrome c.¹³⁹ Apaf-1-deficient fibroblasts were resistant to ceramide-induced apoptosis,¹⁴⁰ and exogenous ceramide-c2-induced apoptosis was inhibited by overexpression of Bcl-2 and Bcl-xL.^{139,141} Ceramide directly induced mitochondrial depolarization and cytochrome c release in isolated mitochondria in some experiments but failed in others. Others found that ceramide was metabolized to disialoganglioside (GD3) in the Golgi apparatus, and caspase dependent accumulation of GD3 was involved in Fas-induced mitochondrial membrane depolarization and apoptosis.¹⁴² Further, ceramide may inhibit protein kinase Akt responsible for the phosphorylation of Bad, which may in turn inhibit Bcl-2.¹⁴³

Since the cytotoxic action of many chemotherapeutic agents is dependent on mitochondrial membrane depolarization and inhibited by Bcl-2, this molecular pathway activated by death receptors could sensitize cells to chemotherapy.

Stress signals induced by chemotherapeutic drugs and DNA damage can induce ceramide synthase activation in the mitochondria and endoplasmic reticulum, probably by mechanisms distinct from Smase activation by death receptors. Activation of ceramide synthase was demonstrated in daunorubicin-induced apoptosis in p388 and U937 cells¹²⁷ and also in HL60 cells.¹⁴¹ The exact mechanism of how DNA damage may induce ceramide synthesis is not known. However, the role of p53 activation has been indicated.¹⁴⁵

Upregulation of FasL in response to chemotherapeutic agents can be dependent on ceramide production and impaired in cells from patients with Niemann-Pick disease deficient in Smase.¹⁰⁴ However ceramide can induce the activation of JNK that leads to activation of AP-1, which in turn can upregulate several death ligands including FasL.⁷⁵ In fact, ceramide-induced apoptosis in HL-60 cells was dependent on AP-1 activation and c-jun expression.¹⁴⁶

Of interest was the report in which exogenous ceramide-C2 induced apoptosis only in type II cells (Fas-mediated apoptosis), such as Jurkat and CEM, but type I cells (H9, SKW6) were not sensitive. Moreover, protein kinase C activation inhibited both Fas- and ceramide- induced apoptosis only in type II cells.¹⁴⁷ Protein kinase C can

induce the activation of sphingosine kinase, the enzyme responsible for the conversion of sphingosine into sphingosine-1-phosphate, a metabolite known to inhibit ceramide-induced apoptosis.¹⁴⁸ Based on these observations we can hypothesize that in type II cells induction of ceramide synthesis by chemotherapeutic agents upregulates FasL and other death ligands leading to activation of death receptors, but at the same time ceramide can directly induce apoptosis via mitochondrial membrane depolarization. In contrast in type I cells, drug-induced apoptosis may be more dependent on the activation of death receptor-induced apoptosis via direct activation of effector caspases independently from the mitochondria. Thus the same chemotherapeutic drug can induce apoptosis dependent on DR-DL interactions in type I cells while being „independent“ in type II cells⁷⁰

Conclusions

Death ligands upregulated during cytotoxic stress in turn can activate an autocrine or paracrine cell death in constitutively sensitive cells. Further, cancer therapy can sensitize resistant tumors to death ligands by upregulating death receptors or via other mechanisms that converge on the pathway leading to apoptosis. Therefore death ligands can be considered as “death genes” since their de novo expression is required for apoptosis induction. It is evident that drug-induced apoptosis can require functional death receptors for the apoptotic signal to be transmitted, however this apoptosis can also proceed independent of Fas. The cell type, the molecular target and the drug concentration determine whether the upregulation of FasL and activation of the Fas pathway is involved in drug-induced apoptosis. Since common pathways between drug-induced and death receptor-induced apoptosis involve direct activation of initiator caspase-8 or of the effector caspases downstream of the mitochondria, aberrant expression of a specific caspase would lead to cross resistance to the two forms of cell death. However cytotoxic stress can directly target the mitochondria, thereby simply bypassing the Fas signal. Drugs that generate ROS or induce ceramide production (e.g. doxorubicin, bleomycin) can rapidly activate the JNK/SAPK pathway and upregulate FasL therefore can induce a Fas-FasL dependent early apoptosis. However, in these model systems Fas dependence is only transient and cell death occurs later induced by alternative signaling mechanisms.

The concept of employing therapeutic strategies to modulate death receptor signaling pathways in combination with anticancer agents is only in its infancy. One approach has involved modulation of the Fas death receptor to sensitize human colon carcinomas to FUra/LV-induced cytotoxicity. Apoptosis downstream of DNA damage induced by FUra/LV is dependent on Fas/FasL interactions and on a

functional Fas signaling pathway. Cells are sensitized to FUra/LV by IFN- γ -induced upregulation of Fas expression.⁷² In human colon carcinoma cells the dependency of FUra/LV-induced thymineless death on Fas signaling is unique to this cytotoxic agent and has not been demonstrated for other DNA damaging agents. Hence the combination of FUra/LV and IFN- γ is completing Phase I clinical trial in patients with colon carcinoma and will subsequently be evaluated as a therapeutic modality in previously untreated patients. This is an excellent example of how therapeutic strategies can be developed to exploit the dependency and crosstalk between death receptor signaling and drug signaling pathways.

Further, even if drug induced cytotoxicity can induce "Fas-independent" apoptosis, the reactivation of death receptor pathways may open alternative cell death mechanisms. Also, the role of death receptors and ligands in the anti-tumor activity of chemotherapeutic agents is even more complex in vivo. Death ligands may kill bystander drug-resistant cells and attract immune competent cells, and upregulation of death receptors can sensitize tumor cells to the immune response.

For the cytotoxic ligand TRAIL, anticancer agents with diverse mechanisms of action including VP-16, doxorubicin, cisplatin, FUra and Ara-C, have demonstrated the ability to enhance TRAIL-induced apoptosis.^{100,101} These agents can upregulate the expression of TRAIL receptors. While the signaling pathway(s) for TRAIL-induced apoptosis remain to be clearly defined, it is probable that the TRAIL signaling and drug signaling pathways converge at various levels in their respective pathways. Therefore the approaches to combine cytokines or ligands that engage death receptor signaling pathways with cytotoxic agents are yielding considerable potential in the exploration of new targets for therapeutic exploitation in the treatment of human cancers.

References

- ^{1.} *Ashkenazi A, Dixit VM*: Death receptors: Signaling and modulation. *Science* 281:305-108, 1998.
- ^{2.} *Ware CF, VanArsdale S, VanArsdale TL*: Apoptosis mediated by the TNF-related cytokine and receptor families. *J Cell Biochem* 60:47-55, 1996.
- ^{3.} *McGeehan GM, Becherer JD, Bast RC Jr, et al*: Regulation of tumor necrosis factor- α processing by a metalloproteinase inhibitor. *Nature* 370:558-561, 1994.
- ^{4.} *Tanaka M, Itai T, Adachi M, et al*: Downregulation of Fas ligand by shedding. *Nature Med* 4:31-36, 1998.
- ^{5.} *Kayagaki N, Kawasaki A, Ebata T, et al*: Metalloproteinase-mediated release of human Fas ligand. *J Exp Med* 182:1777-1783, 1995.
- ^{6.} *Ashkenazi A, Dixit VM*: Apoptosis control by death and decoy receptors. *Curr Opin Cell Biol* 11:255-260, 1999.
- ^{7.} *Nagata S, Golstein P*: The Fas death factor. *Science* 276:1449, 1995.
- ^{8.} *Arai H, Gordon D, Nabel EG, et al*: Gene transfer of Fas ligand induces tumor regression in vivo. *Proc Natl Acad Sci USA* 94:13862-13867, 1997.
- ^{9.} *French LE, Hahne M, Viard I, et al*: Fas and Fas ligand in embryos and adult mice: ligand expression in several immune-privileged tissues and coexpression in adult tissues characterized by apoptotic cell turnover. *J Cell Biol* 133:335-343, 1996.
- ^{10.} *Moller P, Koretz K, Leithauser F, et al*: Expression of Apo-1 (CD95), a member of the NGF/TNF receptor superfamily, in normal and neoplastic colon epithelium. *Int J Cancer* 57:371-377, 1994.
- ^{11.} *Pitti RM, Marsters SA, Lawrence DA, et al*: Genomic amplification of a decoy receptor for Fas ligand in lung and colon cancer. *Nature*. 396:699-703, 1998.
- ^{12.} *Itoh N, Yonehara S, Ishii A, et al*: The polypeptide encoded by the cDNA for human cell surface antigen Fas can mediate apoptosis. *Cell* 66:233-243, 1991.
- ^{13.} *Boldin MP, Varfolomeev EE, Panczer Z, et al*: A novel protein that interacts with the death domain of Fas/APO1 contains a sequence motif related to the death domain. *J Biol Chem* 270:7795-7798, 1995.
- ^{14.} *Chinnaiyan AM, O'Rourke K, Tewari M, et al*: FADD, a novel death domain-containing protein, interacts with the death domain of Fas and initiates apoptosis. *Cell* 81:505-512, 1995.
- ^{15.} *Boldin MP, Goncharov TM, Goltsev YV, et al*: Involvement of MACH, a novel MORT1/FADD-interacting protease, in Fas/APO1 and TNF receptor-induced cell death. *Cell* 85:803-815, 1996.
- ^{16.} *Medema JP, Scaffidi C, Kischkel FC, et al*: FLICE is activated by association with the CD95 death-inducing signaling complex (DISC). *EMBO J* 16:2794-2804, 1997.
- ^{17.} *Muzio M, Chinnaiyan AM, Kischkel FC, et al*: FLICE a novel FADD-homologous ICE/CED-3-like protease is recruited to the CD95 (Fas/Apo-1) death-inducing signaling complex. *Cell* 85:817-827, 1996.
- ^{18.} *Irmeler M, Thome M, Hahne M, et al*: Inhibition of death receptor signals by cellular FLIP. *Nature* 388:190-195, 1997.
- ^{19.} *Scaffidi C, Schmitz I, Krammer PH, et al*: The role of c-FLIP in modulation of CD95-induced apoptosis. *J Biol Chem* 274:1541-1548, 1999.
- ^{20.} *Scaffidi C, Fulda S, Srinivasan A, et al*: Two CD95 (APO-1/Fas) signaling pathways. *EMBO J* 17:1675-1687, 1998.
- ^{21.} *Sun X-M, MacFarlane M, Zhuang J, et al*: Distinct caspase cascades are initiated in receptor-mediated and chemical-induced apoptosis. *J Biol Chem* 274:5053-5060, 1999.
- ^{22.} *Kuwana T, Smith JJ, Muzio M, et al*: Apoptosis induction by caspase-8 is amplified through the mitochondrial release of cytochrome c. *J Biol Chem*. 27:16589-16594, 1998.
- ^{23.} *Li P, Nijhawan D, Alnemri ES, Wang X*: Cytochrome c and dATP-dependent formation of Apaf-1/caspase-9 complex initiates an apoptotic protease cascade. *Cell* 91:479-489, 1997.
- ^{24.} *Jeremias I, Herr I, Boehler T, et al*: TRAIL/Apo-2-ligand-induced apoptosis in human T cells. *Eur J Immunol* 28:143-152, 1998.
- ^{25.} *Marsters SA, Pitti RM, Donahue CJ, et al*: Activation of apoptosis by Apo-2 ligand is independent of FADD but blocked by CrmA. *Curr Biol* 6:750-752, 1996.
- ^{26.} *Griffith TS, Lynch DH*: TRAIL: a molecule with multiple receptors and control mechanisms. *Curr Opin Immunol* 10:559-563, 1998.
- ^{27.} *Zhang XD, Franco AV, Nguyen T, et al*: Differential Localization and Regulation of Death and Decoy Receptors for TNF-Related Apoptosis-Inducing Ligand (TRAIL) in Human Melanoma Cells. *J Immunol* 164:3961-3970, 2000.
- ^{28.} *Walczak H, Miller RE, Ariail K, et al*: Tumoricidal activity of tumor necrosis factor-related apoptosis-inducing ligand in vivo. *Nature Med* 5:157-163, 1999.

- 29.² *Walczak H, Degli-Eposti MA, Johnson RS, et al*: TRAIL-R2: a novel apoptosis-mediating receptor for TRAIL. *EMBO J* 16:5386-5397, 1997.
- 30.² *Wiley SR, Schooley K, Smolak PJ, et al*: Identification and characterization of a new member of the TNF family that induces apoptosis. *Immunity* 3:673-682, 1995.
- 31.² *Pan G, O'Rourke K, Chinnaiyan AM, et al*: The receptor for the cytotoxic ligand TRAIL. *Science* 276:111-113, 1997.
- 32.² *Pan G, Ni J, Wei YF, et al*: An antagonist decoy receptor and a death domain-containing receptor for TRAIL. *Science* 277:815-818, 1997.
- 33.² *Sheridan JP, Marsters SA, Pitti PM, et al*: Control of TRAIL-induced apoptosis by a family of signaling and decoy receptors. *Science* 277:818-821, 1997.
- 34.² *Degli-Eposti MA, Smolak PJ, Walczak H, et al*: Cloning and characterization of TRAIL-R3, a novel member of the emerging TRAIL receptor family. *J Exp Med* 186:1165-1170, 1997.
- 35.² *Degli-Eposti MA, Dougall WC, Smolak PJ, et al*: The novel receptor TRAIL-R4 induces NF- κ B and protects against TRAIL-mediated apoptosis, yet retains an incomplete death domain. *Immunity* 7:813-820, 1997.
- 36.² *Marsters SA, Sheridan JP, Pitti RM, et al*: A novel receptor for Apo2L/TRAIL contains a truncated death domain. *Curr Biol* 7:1003-1006, 1997.
- 37.² *Frank S, Kohler U, Schackert G, et al*: Expression of TRAIL and its receptors in human brain tumors. *Biochem Biophys Res Commun* 257:454-459, 1999.
- 38.² *Griffith TS, Chin WA, Jackson GC, et al*: Intracellular regulation of TRAIL-induced apoptosis in human melanoma cells. *J Immunol* 161:2833-2840, 1998.
- 39.² *Thomas WD, Hersey P*: TNF-related apoptosis-inducing ligand (TRAIL) induces apoptosis in Fas ligand-resistant melanoma cells and mediates CD4 T cell killing of target cells. *J Immunol* 161:2195-2200, 1998.
- 40.² *Ashkenazi A, Pai RC, Fong S, et al*: Safety and antitumor activity of recombinant soluble Apo2 ligand. *J Clin Invest* 104:155-162, 1999.
- 41.² *Rieger J, Naumann U, Glaser T, et al*: APO2 ligand: a novel weapon against malignant glioma? *FEBS Lett* 427:124-128, 1998.
- 42.² *Schneider P, Thome M, Burns K, et al*: TRAIL receptors 1 (DR4) and 2 (DR5) signal FADD-dependent apoptosis and activate NF- κ B. *Immunity* 7:831-6, 1997.
- 42.² *Schneider P, Thome T, Burns K, et al*: TRAIL receptors 1 (DR4) and 2 (DR5) signal FADD-dependent apoptosis and activate NF- κ B. *Immunity* 7:831-836, 1997.
- 43.² *Kischkel FC, Lawrence DA, Chuntharapai A, et al*: Apo2L/TRAIL-dependent recruitment of endogenous FADD and caspase-8 to death receptors 4 and 5. *Immunity* 12:611-620, 2000.
- 44.² *Sprick MR, Weigand MA, Rieser E, et al*: FADD/MORT1 and caspase-8 are recruited to TRAIL receptors 1 and 2 and are essential for apoptosis mediated by TRAIL receptor 2. *Immunity* 12:599-609, 2000.
- 45.² *Bodmer JL, Holler N, Reynard S, et al*: TRAIL receptor-2 signals apoptosis through FADD and caspase-8. *Nat Cell Biol* 2:241-243, 2000.
- 46.² *Hickman JA*: Apoptosis induced by anticancer drugs. *Cancer Metast Rev* 11:121-139, 1992.
- 47.² *Holzman D*: Apoptosis provides new targets for chemotherapy. *J Natl Cancer Inst* 88:1098-1100, 1996.
- 48.² *Sellers WR, Fisher DE*: Apoptosis and cancer drug targeting. *J Clin Invest* 104:1655-1661, 1999.
- 49.² *Miyashita T, Reed JC*: Bcl-2 oncoprotein blocks chemotherapy-induced apoptosis in a human leukemia cell line. *Blood* 81:151-157, 1993.
- 50.² *Bunz F, Hwang PM, Torrance C, et al*: Disruption of p53 in human cancer cells alters the responses to therapeutic agents. *J Clin Invest* 104:263-269, 1999.
- 51.² *Zhu H, Fearnhead HO, Cohen GM*: An ICE-like protease is a common mediator of apoptosis induced by diverse stimuli in human monocytic THP.1 cells. *FEBS LETT* 375:303-308, 1995.
- 52.² *Petak I, Mihalik R, Bauer PI, et al*: BCNU is a caspase-mediated inhibitor of drug-induced apoptosis. *Cancer Res* 58:614-618, 1998.
- 53.² *Petak I, Tillman DM, Harwood FG, et al*: Fas-dependent and -independent mechanisms of cell death following DNA damage in human colon carcinoma cells. *Cancer Res* 60:2643-2650, 2000.
- 54.² *Tounekti O, Pron G, Belehradec J, et al*: Bleomycin, an apoptosis-mimetic drug that induces two types of cell death depending on the number of molecules internalized. *Cancer Res* 53:5462-5469, 1993.
- 55.² *Skladanowski A, Konopa, J*: Adriamycin and daunomycin induce programmed cell death (apoptosis) in tumour cells. *Biochem Pharmacol* 46:375-382, 1993.
- 56.² *Pellicciari C, Bottone MG, Schaack V, et al*: Etoposide at different concentrations may open different apoptotic pathways in thymocytes. *Eur J Histochem* 40:289-298, 1996.
- 57.² *Lock RB, Stribinskiene L*: Dual modes of death induced by etoposide in human epithelial tumor cells allow Bcl-2 to inhibit apoptosis without affecting clonogenic survival. *Cancer Res* 56:4006-4012, 1996.
- 58.² *Kasibhatla S, Brunner T, Genestier L, et al*: DNA damaging agents induce expression of Fas ligand and subsequent apoptosis in T lymphocytes via the activation of NF- κ B and AP-1. *Mol Cell* 1, 543-551, 1998.
- 59.² *Faris M, Kokot N, Latinis K, et al*: The c-Jun N-terminal kinase cascade plays a role in stress-induced apoptosis in Jurkat cells by up-regulating Fas ligand expression. *J Immunol* 160:134-144, 1998.
- 60.² *Friesen C, Herr I, Krammer PH, et al*: Involvement of the CD95 (APO-1/Fas) receptor/ligand system in drug-induced apoptosis in leukemia cells. *Nature Med* 2:574-577, 1996.
- 61.² *Fulda S, Sievens H, Friesen C, et al*: The CD95 (APO1-Fas) system mediates drug-induced apoptosis in neuroblastoma cells. *Cancer Res* 57:3823-3829, 1997.
- 62.² *Villunger A, Egle A, Kos M, et al*: Drug-induced apoptosis is associated with enhanced Fas (APO-1/CD95) ligand expression but occurs independently of Fas (APO-1/CD95) signaling in human T-acute lymphatic leukemia cells. *Cancer Res* 57:3331-3334, 1997.
- 63.² *Eischen CM, Kottke TJ, Martins LM, et al*: Comparison of Apoptosis in wild-type and Fas-resistant cells: chemotherapy-induced apoptosis is not dependent on Fas/Fas ligand interactions. *Blood* 90:935-943, 1997.
- 64.² *Wen J, Ramadevi N, Nguyen D, et al*: Antileukemic drugs increase death receptor 5 levels and enhance apo-2L-induced apoptosis of human acute leukemia cells. *Blood* 96:3900-3906, 2000.
- 65.² *Nagane M, Pan G, Weddle JJ, et al*: Increased death receptor 5 expression by chemotherapeutic agents in human gliomas causes synergistic cytotoxicity with tumor necrosis factor-related apoptosis-inducing ligand in vitro and in vivo. *Cancer Res* 60:847-853, 2000.
- 66.² *Keane MM, Ettenberg SA, Nau MM, et al*: Chemotherapy augments TRAIL-induced apoptosis in breast cell lines. *Cancer Res* 59:734-741, 1999.

- 67.² Alderson MR, Tough TW, Davissmith T, et al: Fas ligand mediates activation-induced cell death in human T lymphocytes. *J Exp Med* 181:71-77, 1995.
- 68.² Brunner T, Mogil RJ, LaFace D, et al: Cell-autonomous Fas (CD95)/Fas-ligand interaction mediates activation-induced apoptosis in T-cell hybridomas. *Nature* 373:441-444, 1995.
- 69.² Fulda S, Strauss G, Meyer E, et al: Functional CD95 ligand and CD95 death-inducing signaling complex in activation-induced cell death and doxorubicin-induced apoptosis in leukemic T cells. *Blood* 95:301-308, 2000.
- 70.² Fulda S, Meyer E, Friesen C, et al: Cell type specific involvement of death receptor and mitochondrial pathways in drug-induced apoptosis. *Oncogene* 20:1063-1075, 2001.
- 71.² Houghton JA, Harwood FG, Tillman DM: Thymineless death in colon carcinoma cells is mediated via Fas signaling. *Proc Natl Acad Sci USA* 94:8144-8149, 1997.
- 72.² Tillman DM, Petak I, Houghton JA: A Fas-dependent component in 5-fluorouracil/leucovorin-induced cytotoxicity in colon carcinoma cells. *Clin Cancer Res* 5:425-430, 1999.
- 73.² Eichhorst ST, Muller M, Li-Weber M, et al: A novel AP-1 element in the CD95 ligand promoter is required for induction of apoptosis in hepatocellular carcinoma cells upon treatment with anticancer drugs. *Mol Cell Biol* 20:7826-7837, 2000.
- 74.² Eichhorst ST, Muerkoster S, Weigand MA, et al: The chemotherapeutic drug 5-fluorouracil induces apoptosis in mouse thymocytes in vivo via activation of the CD95(APO-1/Fas) system. *Cancer Res* 61:243-248, 2001.
- 75.² Harwood FG, Kasibhatla S, Petak I, et al: Regulation of FasL by NF-kappaB and AP-1 in Fas-dependent thymineless death of human colon carcinoma cells. *J Biol Chem* 275:10023-10029, 2000.
- 76.² Muller M, Wilder S, Bannasch D, et al: P53 activates the CD95 (APO-1/Fas) gene in response to DNA damage by anticancer agents. *J Exp Med.*, 188:2033-2045, 1998.
- 77.² Petak I, Tillman DM, Houghton JA: P53-dependence of Fas induction and acute apoptosis in response to 5-fluorouracil-leucovorin in human colon carcinoma cell lines. *Clin Cancer Res* 6:4432-4441, 2000.
- 78.² Siltanen T, Mantymaa P, Saily M, et al: Etoposide-induced apoptosis is not associated with the fas pathway in acute myeloblastic leukemia cells. *Leukemia Res* 24:281-288, 2000.
- 79.² Tolomeo M, Dusonchet L, Meli M, et al: The CD95/CD95 ligand system is not the major effector in anticancer drug-mediated apoptosis. *Cell Death Diff* 5:735-742, 1998.
- 80.² Shao R-G, Cao C-X, Nieves-Neira W et al: Activation of the Fas pathway independently of Fas ligand during apoptosis induced by camptothecin in p53 mutant colon carcinoma cells. *Oncogene* 20:1852-1859, 2001.
- 81.² Wieder T, Essmann F, Prokop A, et al: Activation of caspase-8 in drug-induced apoptosis of B-lymphoid cells is independent of CD95/Fas receptor-ligand interaction and occurs downstream of caspase-3. *Blood* 97:1378-1387, 2001.
- 82.² Wesselborg S, Engels IH, Rossmann E, et al: Anticancer drugs induce caspase-8/FLICE activation and apoptosis in the absence of CD95 receptor/ligand interaction. *Blood* 93:3053-3063, 1999.
- 83.² Tang D, Lahti JM, Kidd VJ: Caspase-8 activation and Bid cleavage contribute to MCF7 cellular execution in a caspase-3-dependent manner during staurosporine-mediated apoptosis. *J Biol Chem* 275:9303-9307, 2000.
- 84.² Ferreira CG, Span SW, Peters GJ: Chemotherapy triggers apoptosis in a caspase-8-dependent and mitochondria-controlled manner in the non-small cell lung cancer cell line NCI-H460. *Cancer Res* 60:7133-7141, 2000.
- 85.² Kinoshita H, Yoshikawa H, Shiiki K, et al: Cisplatin (CDDP) sensitizes human osteosarcoma cells to Fas/CD95-mediated apoptosis by downregulating FLIP-L expression. *In J Cancer* 88:986-991, 2000.
- 86.² Tang D, Lahti JM, Grenet J, et al: Cycloheximide-induced T-cell death is mediated by a Fas-associated death domain-dependent mechanism. *J Biol Chem* 274:7245-7252, 1999.
- 87.² Dorrie J, Schuh W, Keil A, et al: Regulation of CD95 expression and CD95-mediated cell death by interferon-gamma in acute lymphoblastic leukemia with chromosomal translocation t(4;11). *Leukemia* 13:1539-1547, 1999.
- 88.² Griffith TS, Chin WA, Jackson GC, et al: Intracellular regulation of TRAIL-induced apoptosis in human melanoma cells. *J Immunol* 161:2833-2840, 1998.
- 89.² Bretz JD, Rymaszewski M, Arscott PL, et al: TRAIL death pathway expression and induction in thyroid follicular cells. *J Biol Chem* 274:23627-23632, 1999.
- 90.² Kataoka T, Schroter M, Hahne M, et al: FLIP prevents apoptosis induced by death receptors but not by perforin/granzyme B, chemotherapeutic drugs, and gamma radiation. *J Immunol* 161:3936-3942, 1998.
- 91.² Los M, Herr I, Friesen C, et al: Cross-resistance of CD95- and drug-induced apoptosis as a consequence of deficient activation of caspases (ICE/Ced-3 proteases). *Blood* 90:3118-3129, 1997.
- 92.² Boesen-de Cock JG, Tepper AD, de Vries E, et al: Common regulation of apoptosis signaling induced by CD95 and the DNA-damaging stimuli etoposide and -radiation downstream from caspase-8 activation. *J Biol Chem* 274:14255-14261, 1999.
- 93.² Sun X-M, MacFarlane M, Zhuang J, et al: Distinct caspase cascades are initiated in receptor-mediated and chemical-induced apoptosis. *J Biol Chem* 274:50530-50560, 1999.
- 94.² Wu X-X, Mizutani Y, Kakehi Y, et al: Enhancement of Fas-mediated apoptosis in renal cell carcinoma cells by adriamycin. *Cancer Res* 60:2912-2918, 2000.
- 95.² Tamura T, Aoyama N, Saya H, et al: Induction of Fas-mediated apoptosis in p53-transfected human colon carcinoma cells. *Oncogene*, 11:1939-1946, 1995.
- 96.² Owen-Sachau LB, Zhang W, Cusack JC, et al: Wild-type human p53 and a temperature-sensitive mutant induce Fas/APO-1 expression. *Mol Cell Biol* 15:3032-3040, 1995.
- 97.² Rakkar ANS, Katayos Y, Kim M, et al: A novel adenoviral vector expressing human Fas/CD95/APO-1 enhances p53-mediated apoptosis. *Cell Death Diff* 6:326-333, 1999.
- 98.² Fukazawa T, Fujiwara T, Morimoto Y, et al: Differential involvement of the CD95 (Fas/APO-1) receptor/ligand system on apoptosis induced by the wild-type p53 gene transfer in human cancer cells. *Oncogene*, 18:2189-2199, 1999.
- 99.² Shimizu M, Yoshimoto T, Nagata S, et al: A trial to kill tumor cells through Fas (CD95)-mediated apoptosis in vivo. *Biochem Biophys Res Commun* 228:375-379, 1996.
- 100.² Wu GS, Burns TF, McDonald ER 3rd, et al: KILLER/DR5 is a DNA damage-inducible p53-regulated death receptor gene. *Nat Genet* 17:141-143, 1997.
- 101.² Nagane M, Pan G, Weddle JJ, et al: Increased death receptor 5 expression by chemotherapeutic agents in human gliomas causes synergistic cytotoxicity with tumor necrosis factor-related apoptosis-inducing ligand in vitro and in vivo. *Cancer Res* 60:847-853, 2000.
- 102.² Herr I, Wilhelm D, Bohler T, et al: JNK/SAPK activity is not sufficient for anticancer therapy-induced apoptosis involving CD95-L, TRAIL and TNF-alpha. *Int J Cancer* 80:417-424, 1999.

- 103.² *Friesen C, Fulda S, Debatin KM*: Induction of CD95 ligand and apoptosis by doxorubicin is modulated by the redox state in chemosensitive- and drug-resistant tumor cells. *Cell Death Differ* 6:471-480, 1999.
- 104.² *Herr I, Wilhelm D, Bohler T, et al*: Activation of CD95 (APO-1/Fas) signaling by ceramide mediates cancer therapy-induced apoptosis. *EMBO J* 16:6200-6208, 1997.
- 105.² *Boland MP, Foster SJ, O'Neill LAJ*: Daunorubicin activates NF- κ B and induces κ B-dependent gene expression in HL-60 promyelocytic and Jurkat T lymphoma cells. *J Biol Chem* 272:12952-12960, 1997
- 106.² *Miyamoto S, Verma IM*: Rel/NF-kappaB/IkappaB story. *Adv Cancer Res* 66:255-292, 1995.
- 107.² *Baldwin AS*: The NF-kappaB and I kappaB proteins: new discoveries and insights. *Ann Rev Immunol* 14:649-683, 1996.
- 108.² *Chen Z, Hagler J, Palombella VJ, et al*: Signal-induced site-specific phosphorylation targets I kappa B alpha to the ubiquitin-proteasome pathway. *Genes and Dev* 9:1586-1597, 1995.
- 109.² *Karin M*: The regulation of AP-1 activity by mitogen-activated protein kinases. *J Biol Chem* 270:16483-16486, 1995.
- 110.² *Kyriakis J, Banerjee P, Nikolakaki E, et al*: The stress-activated protein kinase subfamily of c-Jun kinases. *Nature* 369:156-160, 1994.
- 111.² *Minden A, Lin A, McMahon M, et al*: Differential activation of ERK and JNK mitogen-activated protein kinases by Raf-1 and MEKK. *Science* 266:1719-1723, 1994.
- 112.² *Derijard B, Hibi M, Wu I-H, et al*: JNK1: a protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell* 76:1025-1037, 1994.
113. *Chen YR, Wang X, Templeton D, et al*: The role of c-Jun N-terminal kinase (JNK) in apoptosis induced by ultraviolet C and radiation. *J Biol Chem* 271:31929-31936, 1996.
- 114.² *Kharbanda S, Ren R, Pandey P, et al*: Activation of the c-Abl tyrosine kinase in the stress response to DNA-damaging agents. *Nature* 376:785-788, 1995.
- 115.² *Van Dam H, Wilhelm D, Herr I, et al*: ATF-2 is preferentially activated by stress-activated protein kinases to mediate c-jun induction in response to genotoxic agents. *EMBO J* 14:1798-1811, 1995.
- 116.² *Yu R, Shtil AA, Tan FH, et al*: Adriamycin activates c-jun N-terminal kinase in human leukemia cells: a relevance to apoptosis. *Cancer Lett* 107:73-81, 1996.
- 117.² *Seimiya H, Mashima T, Toho M, et al*: c-Jun NH2-terminal kinase-mediated activation of interleukin-1 β converting enzyme/CED-3-like protease during anticancer drug-induced apoptosis. *J Biol Chem* 272:4631-4636, 1997.
- 118.² *Sanchez-Perez I, Murguía JR, Perona R*: Cisplatin induces a persistent activation of JNK that is related to cell death. *Oncogene* 16:533-540, 1998.
- 119.² *Rudel T, Zenke FT, Chuang TH, et al*: Cutting edge: p21-activated kinase (PAK) is required for Fas-induced JNK activation in Jurkat cells. *J Immunol* 160:7-11, 1998.
- 120.² *Liu Z-G, Baskaran C, Lee-Choe ET, et al*: Three distinct signaling responses by murine fibroblasts to genotoxic stress. *Nature* 384:273-276, 1996.
- 121.² *Enari M, Sakahiera H, Yokoyama H, et al*: A caspase-activated Dnase that degrades DNA during apoptosis, and its inhibitor ICAD. *Nature* 391:43, 1998.
- 122.² *Lee FS, Hagler J, Chen CJ, et al*: Activation of the I κ B κ kinase complex by MEKK1, a kinase of the JNK pathway. *Cell* 88:213-222, 1997.
- 123.² *Bessho R, Matsubara K, Kubota M, et al*: Pyrrolidine dithiocarbamate, a potent inhibitor of nuclear factor κ B (NF- κ B) activation, prevents apoptosis in human promyelocytic leukemia HL-60 cells and thymocytes. *Biochem Pharmacol* 48:1883-1889, 1994.
- 124.² *Piret B, Piette J*: Topoisomerase poisons activate the transcription factor NF- κ B in ACH-2 and CEM cells. *ucl Acids Res* 24:4242-4248, 1996.
- 125.² *Brach MA, Kharbanda SM, Herrmann F, et al*: Activation of the transcription factor kappaB in human KG-1 myeloid leukemia cells treated with 1-beta-D-arabinofuranosylcytosine. *Mol Pharmacol* 41:60-63, 1992.
- 126.² *Testi R*: Sphingomyelin breakdown and cell fate. *Trends Biochem Sci* 21:468-471, 1996.
- 127.² *Bose R, Verheij M, Haimovitz-Friedman A, et al*: Ceramide synthase mediates daunorubicin-induced apoptosis: an alternative mechanism for generating death signals. *Cell* 82:405-414, 1995.
- 128.² *Hannun YA*: Functions of ceramide in coordinating cellular responses to stress. *Science* 274:1855-1859, 1996.
- 129.² *Hannun YA, Obeid LM*: Mechanisms of ceramide-mediated apoptosis. *Adv Exp Med Biol* 407:145-149, 1997.
- 130.² *Mathias S, Pena LA, Kolesnick RN*: Signal transduction of stress via ceramide. *Biochem J* 335:465-480, 1998.
- 131.² *Verheij M, Bose R, Lin XH, et al*: Requirement for ceramide-initiated SAPK/JNK signalling in stress-induced apoptosis. *Nature* 380:75-79, 1996.
- 132.² *Wiegmann K, Schutze S, Machleidt T, et al*: Functional dichotomy of neutral and acidic sphingomyelinases in tumor necrosis factor signaling. *Cell* 78:1005-1015, 1994.
- 133.² *Spiegel S, Foster D, Kolesnick R*: Signal transduction through lipid second messengers. *Curr Opin Cell Biol* 8:159-167, 1996.
- 134.² *Hirschberg K, Rodger J, Futerman AH*: The long-chain sphingoid base of sphingolipids is acylated at the cytosolic surface of the endoplasmic reticulum in rat liver. *Biochem J* 290:751-757, 1993.
- 135.² *Shimeno H, Soeda S, Yasukouchi M, et al*: Fatty acyl-Co A: sphingosine acyltransferase in bovine brain mitochondria: its solubilization and reconstitution onto the membrane lipid liposomes. *Biol Pharm Bull* 18:1335-1339, 1995.
- 136.² *Adam-Klages S, Schwandner R, Adam D, et al*: Distinct adapter proteins mediate acid versus neutral sphingomyelinase activation through the p55 receptor for tumor necrosis factor. *J Leukoc Biol* 63:678-682, 1998.
- 137.² *Schwandner R, Wiegmann K, Bernardo K, et al*: TNF receptor death domain-associated proteins TRADD and FADD signal activation of acid sphingomyelinase. *J Biol Chem* 273:5916-5922, 1998.
- 138.² *Wiegmann K, Schwandne R, Krut O, et al*: Requirement of FADD for tumor necrosis factor-induced activation of acid sphingomyelinase. *J Biol Chem* 274:5267-5270, 1999.
- 139.² *Cuvillier O, Edsall L, Spiegel S*: Involvement of sphingosine in mitochondria-dependent Fas-induced apoptosis of type II Jurkat T cells. *J Biol Chem* 275:15691-15700, 2000.
- 140.² *Cecconi F, Alvarez-Bolado G, Meyer BI, et al*: Apaf1 (CED-4 homolog) regulates programmed cell death in mammalian development. *Cell* 94:727-737, 1998.
- 141.² *Susin SA, Zamzami N, Castedo M, et al*: The central executioner of apoptosis: multiple connections between protease activation and mitochondria in Fas/APO-1/CD95- and ceramide-induced apoptosis. *J Exp Med* 186:25-37, 1997.
- 142.² *De Maria R, Lenti L, Malisan F, et al*: Requirement for GD3 ganglioside in CD95- and ceramide-induced apoptosis. *Science* 277:1652-1655, 1997.
- 143.² *Zhou H, Summers SA, Birnbaum MJ, et al*: Inhibition of Akt kinase by cell-permeable ceramide and its implications for

- ceramide-induced apoptosis. *J Biol Chem* 273:16568-16575, 1998.
- 144.²*Boland MP, Foster SJ, O'Neill LA*: Daunorubicin activates NFkappaB and induces kappaB-dependent gene expression in HL-60 promyelocytic and Jurkat T lymphoma cells. *J Biol Chem* 272:12952-12960, 1997.
- 145.²*Dbaiho GS, Pushkareva MY, Rachid RA, et al*: p53-dependent ceramide response to genotoxic stress. *J Clin Invest* 102:329-339, 1998.
- 146.²*Sawai H, Okazaki T, Yamamoto H, et al*: Requirement of AP-1 for ceramide-induced apoptosis in human leukemia HL-60 cells. *J Biol Chem* 270:27326-27331, 1995.
- 147.²*Scaffidi C, Schmitz I, Zha J, et al*: Differential modulation of apoptosis sensitivity in CD95 type I and type II cells. *J Biol Chem* 274:22532-22538, 1999.
- 148.²*Cuvillier O, Pirianov G, Kleuser B, et al*: Suppression of ceramide-mediated programmed cell death by sphingosine-1-phosphate. *Nature* 381:800-803, 1996.